

Notes and Comments

Length at first sexual maturity of the freshwater fish fauna of the Baixada Maranhense Environmental Protection Area

Y. B. S. Nunes^{a*} , R. Milke^b , L. R. Silva^b , C. L. França^c , C. E. S. S. Costa^d , J. L. S. Nunes^e  and M. B. Figueiredo^b 

^aUniversidade Federal Rural da Amazônia – UFRA, Programa de Pós-graduação em Aquicultura e Recursos Aquáticos Tropicais, Belém, PA, Brasil

^bUniversidade Estadual do Maranhão – UEMA, Programa de Pós-graduação em Ecologia e Conservação da Biodiversidade, Laboratório de Biologia Pesqueira, São Luís, MA, Brasil

^cUniversidade Federal do Rio Grande – FURG, Programa de Pós-graduação em Aquicultura, Estação Marinha de Aquicultura, Rio Grande, RS, Brasil

^dUniversidade Estadual do Maranhão – UEMA, Departamento de Engenharia de Pesca, São Luís, MA, Brasil

^eUniversidade Federal do Maranhão – UFMA, Departamento de Oceanografia e Limnologia, São Luís, MA, Brasil

Despite ongoing research efforts, the ichthyofauna of the coastal basins of Maranhão State, Brazil remains poorly described (Abreu et al., 2019). In recent years the number of research publications on fish diversity in this region has increased (e. g. Brito et al., 2019; Guimarães et al., 2020; Oliveira et al., 2020), however important work remains to be done. An accurate assessment of population parameters related to fish reproduction is an essential component of effective fisheries management (Brown-Peterson et al., 2011; Nascimento et al., 2015; Nunes et al., 2019).

The Baixada Maranhense Environmental Protection Area is a RAMSAR site consisting of a complex ecosystem in which human beings play an essential role in managing, using, and conserving several components. Due to the abundance of regional water resources, fishing is among the most important socioeconomic activities (Araujo and Pinheiro, 2008). In recent decades this region has undergone substantial anthropogenic changes, decreased river water flow, siltation, damming, and other anthropic activities compounded by changes in rainfall patterns (Cantanhêde et al., 2017). Monitoring the ichthyofauna in rivers is essential to identify environmental responses to human actions and provides subsidies to regulate the use of water resources, thus enabling the development of alternatives to minimize river degradation (Santos et al., 2017; Lima et al., 2018).

This study describes the minimum size at first maturation (L_{50}) of commercially important fish species in two distinct habitats (a river and a lake) within the Pindaré River Hydrographic Basin. We use data from monthly collections that were carried out for 18 months (January 2015 to June 2016), in Lake Viana and the Pindaré river in the state of Maranhão, in northeastern Brazil. The study area is a designated RAMSAR Site, protected by Brazil under the RAMSAR intergovernmental treaty (Brasil, 2010). It is characterized by seasonal changes in rainfall that influence both the reproductive behavior and adaptations of fish

species and the dynamics of the local fishing industry. When there is a shortage of fish stocks, extractive activity is intensified to meet demand, and the seasonal reduction in floods that occur annually triggers annual overfishing (Carvalho et al., 2017; Carvalho et al., 2021).

We purchased specimens used for this work from local fish markets, then stored them on ice for transport to the laboratory, where they were subsequently identified to the lowest possible taxonomic level. We then analyzed all specimens morphologically to assess total length (TL) and total weight (TW) with the aid of an ichthyometer and an electronic caliper with a precision of 1 mm and 0.01 g, respectively. Then we made a longitudinal incision on the ventral portion of each specimen to remove the gonads. We then fixed the gonads in a 5% formalin solution for subsequent preservation in 70% ethanol. We deposited all specimens in the Collection of tissues and DNA of the fauna of Maranhão, UEMA (CoFauMA), and in the teaching collection of fishes of the Laboratory of Fisheries and Aquatic Ecology, UEMA (LabPEA).

We assessed the reproductive state for each specimen through macroscopic identification of the stage of gonadal maturation and characterization of the gonads according to the scale proposed by Brown-Peterson et al. (2011) and Lowerre-Barbieri et al. (2011). We calculated minimum size at first maturation (L_{50}) based on the cumulative curve of occurrence frequencies of adult individuals by total length class adjusted to the logistic function. We characterized the maturation as A (immature); B (under development); C (able to spawn); D (regressing), and E (regenerated). We calculated the percentage of adult organisms (stages B, C, D, and E) by length class and considered as a dependent variable (Y) and the total length as an independent variable (X). Subsequently, we fitted these values to a logistic curve, using the Statistica 7 Program according to the following formula: $P = 1/(1 + \exp[-r(L - L_m)])$. P is the proportion

*e-mail: ybruno3098@gmail.com

Received: September 24, 2021 – Accepted: December 7, 2021



This is an Open Access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

of mature individuals, **r** is the slope of the curve, **L** is the length, and **L_m** is the average length of sexual maturity.

Overall, we analyzed 1,324 fish (Table 1) representing ten species: *Curimata* sp. (n = 61), *Platydoras brachylecis* Piorski, Garavello, Arce, Sabaj and Pérez, 2008 (n = 97), *Prochilodus lacustris* Steindachner, 1907 (n = 61), *Plagioscion squamosissimus* Heckel, 1840 (n = 205), *Hoplias malabaricus* Bloch, 1784 (n = 94), *Trachelyopterus galeatus* Linnaeus, 1766 (n = 211), *Ageneiosus dentatus* Kner, 1858 (n = 91), *Schizodon dissimilis* Garman, 1890 (n = 252), *Hassar affinis* Steindachner, 1881 (n = 14), and *Pygocentrus nattereri* Kner, 1858 (n = 105).

Our study showed that *P. squamosissimus*, *H. affinis*, *H. malabaricus* and *P. lacustris* had a L₅₀ higher than that calculated in the literature, as shown in Table 2. The increase in the minimum length at first maturity can be an indication

of stock recovery or an increase in the availability of food resources for the species (Santos et al., 2015). On the other hand, the species *S. dissimilis*, *P. nattereri* and *T. galeatus* presented L₅₀ with values lower in this survey (Table 2). According to Camargo and Lima Junior (2007), the decrease in the L₅₀ could mean a potential danger of overfishing. This decrease may indicate that fish are reaching the minimum length of first maturation early as a need for stock renewal (Ikeda, 2003). Finally, the individuals *A. dentatus*, *Curimata* sp., and *P. brachylecis* still do not have literature with L₅₀ records, precluding historical comparisons.

Based on a robust sample size from a previously poorly-characterized ichthyofauna, we showed that the estimates of L₅₀ for several species collected in the Baixada Maranhense Environmental Protection Area differ from those reported in the literature. Although fishing activity

Table 1. Variation in total length and L₅₀ for freshwater fish from northeastern Brazil.

SPECIES	Total length (cm)			L50 considering both sexes (%)	L50 both sexes (cm)	MALE				FEMALE			
	MIN	MAX	± SD			MIN	MAX	Immature (%)	Maturity equation	MIN	MAX	Immature (%)	Maturity equation
<i>Ageneiosus dentatus</i> (Kner, 1858)	13.2	26.7	19.44 ± 2.33	35.16	17.19	14.6	24.0	52.00	$y=1/(1+\exp(-(0.35)^x(x-17.99)))$	13.2	26.7	14.63	$y=1/(1+\exp(-(0.74)^x(x-15.30)))$
<i>Curimata</i> sp.	11.3	49.0	16.72 ± 8.12	16.39	13.98	11.9	33.0	25.00	$y=1/(1+\exp(-(0.62)^x(x-19.61)))$	11.3	49.0	10.81	$y=1/(1+\exp(-(0.51)^x(x-14.83)))$
<i>Hassar affinis</i> (Steindachner, 1881)	11.1	17.0	13.89 ± 1.33	38.1	12.98	11.1	17.0	39.02	$y=1/(1+\exp(-(0.60)^x(x-13.06)))$	11.3	17.0	36.92	$y=1/(1+\exp(-(0.77)^x(x-13.04)))$
<i>Hoplias malabaricus</i> (Bloch, 1784)	17.3	32.1	24.05 ± 2.98	30.85	20.67	18.0	28.2	60.71	$y=1/(1+\exp(-(0.56)^x(x-22.21)))$	17.3	32.1	18.18	$y=1/(1+\exp(-(1.21)^x(x-18.44)))$
<i>Plagioscion squamosissimus</i> (Heckel, 1840)	15.5	39.8	24.03 ± 3.63	36.59	25.32	15.5	39.8	34.31	$y=1/(1+\exp(-(0.25)^x(x-25.16)))$	17.0	34.5	39.71	$y=1/(1+\exp(-(0.30)^x(x-21.15)))$
<i>Platydoras brachylecis</i> (Piorski, Garavello, Arce, Sabaj e Pérez, 2008)	10.5	23.0	14.78 ± 1.88	28.87	13.23	10.5	18.5	56.52	$y=1/(1+\exp(-(0.59)^x(x-14.46)))$	11.7	23.0	3.94	$y=1/(1+\exp(-(2.28)^x(x-10.95)))$
<i>Prochilodus lacustris</i> (Steindachner, 1907)	13.9	21.3	17.54 ± 1.74	4.92	15.07	13.9	21.1	3.23	$y=1/(1+\exp(-(2.10)^x(x-15.01)))$	14.9	21.3	6.67	$y=1/(1+\exp(-(1.57)^x(x-15.16)))$
<i>Pygocentrus nattereri</i> (Kner, 1858)	12.4	19.2	15.39 ± 1.59	31.43	13.9	12.4	18.9	31.37	$y=1/(1+\exp(-(0.61)^x(x-13.88)))$	12.4	19.2	31.48	$y=1/(1+\exp(-(0.96)^x(x-14.05)))$
<i>Schizodon dissimilis</i> (Garman, 1890)	16.5	38.5	23.22 ± 3.49	36.9	20.25	17.4	31.9	40.00	$y=1/(1+\exp(-(0.36)^x(x-20.22)))$	16.5	38.5	34.87	$y=1/(1+\exp(-(0.43)^x(x-20.37)))$
<i>Trachelyopterus galeatus</i> (Linnaeus, 1766)	14	17.4	12.37 ± 1.56	33.65	11.13	14	17.4	36.52	$y=1/(1+\exp(-(0.24)^x(x-11.16)))$	14	14.8	30.21	$y=1/(1+\exp(-(0.44)^x(x-9.53)))$

Caption: MIN = minimum total length (cm); MAX = maximum total length (cm); SD = standard deviation.

Table 2. Differences in estimated L_{50} between the present work and previously published analyses.

Species	Authors	L50 Previous estimate	L50 Present study	Classification
<i>Hassar affinis</i>	Cantanhêde et al. (2016) and Carvalho et al. (2021)	11.52 and 12.3	12.98	Upper
<i>Hoplias malabaricus</i>	Carvalho et al. (2021)	19.0	20.67	Upper
<i>Plagioscion squamosissimus</i>	Carvalho et al. (2017, 2021)	15.3 and 17.3	25.32	Upper
<i>Prochilodus lacustris</i>	Carvalho et al. (2021)	14.4	15.07	Upper
<i>Pygocentrus nattereri</i>	Carvalho et al. (2021)	14.3	13.00	Less
<i>Schizodon dissimilis</i>	Carvalho et al. (2021)	25.0	20.25	Less
<i>Trachelyopterus galeatus</i>	Garcia et al. (2019)	14.5	11.13	Less

characteristic of this RAMSAR site may affect estimates of L_{50} , this study can help to understand the renewal of fish stocks, emphasizing species that do not yet have records of minimum capture size (*Ageneiosus dentatus*, *Curimata* sp. and *Platydoras brachylecis*). Thus, it will serve as a basis for adopting measures that make viable the practice of sustainable fishing, contributing to inspection and adjustment in the sizes of fishing gear.

References

- ABREU, J.M.S., CRAIG, J.M., ALBERT, J.S. and PIORSKI, N.M., 2019. Historical biogeography of fishes from coastal basins of Maranhão State, northeastern Brazil. *Neotropical Ichthyology*, vol. 17, no. 2, pp. e180156. <http://dx.doi.org/10.1590/1982-0224-20180156>.
- ARAUJO, N.A. and PINHEIRO, C.U.B., 2008. Avaliação sócio-econômica da pesca artesanal e do potencial aquícola na região lacustre de Penalva - APA da Baixada Maranhense. *Boletim do Laboratório de Hidrobiologia*, vol. 21, no. 1, pp. 41-50.
- BRASIL. Ministério do Meio Ambiente - MMA, 2010 [viewed 30 December 2010]. *Convenção RAMSAR* [online]. Available from: www.mma.gov.br
- BRITO, P.S., GUIMARÃES, E.C., FERREIRA, B.R.A., OTTONI, F.P. and PIORSKI, N.M., 2019. Freshwater fishes of the Parque Nacional dos Lençóis Maranhenses and adjacent areas. *Biota Neotropica*, vol. 19, no. 3, pp. e20180660. <http://dx.doi.org/10.1590/1676-0611-bn-2018-0660>.
- BROWN-PETERSON, N.J., WYANSKI, D.M., SABORIDO-REY, F., MACEWICZ, B.J. and LOWERRE-BARBIERI, S.K., 2011. A standardized terminology for describing reproductive development in fishes. *Marine and Coastal Fisheries*, vol. 3, no. 1, pp. 52-70. <http://dx.doi.org/10.1080/19425120.2011.555724>.
- CAMARGO, M. and LIMA JÚNIOR, W.M.A., 2007. Aspectos da biologia reprodutiva de seis espécies de peixes de importância comercial do Médio Rio Xingu - Bases para seu manejo. *Scientific Magazine UAKARI*, vol. 3, no. 1, pp. 64-77. <http://dx.doi.org/10.31420/uakari.v3i1.20>.
- CANTANHÊDE, L.G., CARVALHO, I.F.D.S., NUNES, K.B., SANTOS, N.B. and ALMEIDA, Z.D.S.D., 2017. Reproductive ecology of the catfish, *Hassar affinis* (ACTINOPTERYGII: DORADIDAE), in three lakes of the Pindaré-Mearim Lake System, Maranhão. *Ciência Agrônômica*, vol. 48, no. 3, pp. 464-472. <http://dx.doi.org/10.5935/1806-6690.20170054>.
- CANTANHÊDE, L.G., CARVALHO, I.F.S., SANTOS, N.B. and ALMEIDA, Z.S., 2016. Biologia reprodutiva do *Hassar affinis* (Pisces: Siluriformes, Doradidae), Lago de Viana, Baixada Maranhense, Maranhão, Brasil. *Acta Amazonica*, vol. 46, no. 2, pp. 219-226. <http://dx.doi.org/10.1590/1809-4392201503844>.
- CARVALHO, I.F.S., CANTANHÊDE, L.G., DINIZ, A.L.C., CARVALHO-NETA, R.N.F. and ALMEIDA, Z.S., 2021. Reproductive biology of seven fish species of commercial interest at the Ramsar site in the Baixada Maranhense, Legal Amazon, Brazil. *Neotropical Ichthyology*, vol. 19, no. 2, pp. e200067. <http://dx.doi.org/10.1590/1982-0224-2020-0067>.
- CARVALHO, I.F.S., CANTANHÊDE, L.G., SANTOS, N.B., CARVALHO-NETA, R.N.F. and ALMEIDA, Z.S., 2017. Biologia reprodutiva de *Plagioscion squamosissimus* (Pisces, Sciaenidae) em uma área de proteção ambiental do Nordeste do Brasil. *Boletim do Instituto de Pesca*, vol. 43, no. 2, pp. 243-256. <http://dx.doi.org/10.20950/1678-2305.2017v43n2p243>.
- GARCIA, D.A.Z., VIDOTTO-MAGNONI, A.P. and ORSI, M.L., 2019. Características reprodutivas de peixes invasores no rio Paranapanema, bacia do alto rio Paraná, sul do Brasil. *Neotropical Biology and Conservation*, vol. 14, no. 4, pp. 511-528. <http://dx.doi.org/10.3897/neotropical.14.e49079>.
- GUIMARÃES, E.C., BRITO, P.S., GONÇALVES, C. and OTTONI, F.P., 2020. An inventory of Ichthyofauna of the Pindaré River drainage, Mearim River basin, Northeastern Brazil. *Biota Neotropica*, vol. 20, no. 4, pp. e20201023. <http://dx.doi.org/10.1590/1676-0611-bn-2020-1023>.
- IKEDA, R.G.P., 2003. *Idade, crescimento e aspectos reprodutivos de Macrodon ancylodon (Bloch e Schneider, 1801), na Costa Norte do Brasil*. São Paulo: Universidade de São Paulo, 87 p. Dissertação de Mestrado em Ciências.
- LIMA, F.P., NOBILE, A.B., FREITAS-SOUZA, D., CARVALHO, E.D. and VIDOTTO-MAGNONI, A.P., 2018. Can dams affect the trophic structure of ichthyofauna? A long-term effect in the Neotropical region. *Iheringia. Série Zoologia*, vol. 108, no. 0. <http://dx.doi.org/10.1590/1678-4766e2018030>.
- LOWERRE-BARBIERI, S.K., GANIAS, K., SABORIDO-REY, F., MURUA, H. and HUNTER, J.R., 2011. Reproductive timing in marine fishes: variability, temporal scales, and methods. *Marine and Coastal Fisheries*, vol. 3, no. 1, pp. 71-91. <http://dx.doi.org/10.1080/19425120.2011.556932>.

- NASCIMENTO, W.S., YAMAMOTO, M.E., CHELLAPPA, N.T., ROCHA, O. and CHELLAPPA, S., 2015. Conservation status of an endangered annual fish *Hypsolebias antenori* (Rivulidae) from Northeastern Brazil. *Brazilian Journal of Biology = Revista Brasileira de Biologia*, vol. 75, no. 2, pp. 484-490. <http://dx.doi.org/10.1590/1519-6984.19113>. PMID:26132036.
- NUNES, J.L.S., COSTA, M.S., RIBEIRO, M.F.R., COSTA, M.S. and MENDES, M.B.P., 2019. Length weight relationship for freshwater fish species from Brazilian Cerrado. *Journal of Applied Ichthyology*, vol. 35, pp. 1-3. <http://dx.doi.org/10.1111/jai.13889>.
- OLIVEIRA, E.S., GUIMARÃES, E.C., BRITO, P.S., VIEIRA, L.A., OLIVEIRA, R.F., CAMPOS, D.S., KATZ, A.M., SOUTH, J., NUNES, J.L.S. and OTTONI, F.P., 2020. Ichthyofauna of the Mata de Itamaçoca, an urban protected area from the upper Munim River basin, Northern Brazilian Cerrado. *Biota Neotropica*, vol. 20, no. 4, pp. e20201116. <http://dx.doi.org/10.1590/1676-0611-bn-2020-1116>.
- SANTOS, A.C., GONÇALVES, C.C. and CARVALHO, F.R., 2017. Ichthyofauna of the “Cachoeira de São Roberto” and fishes of lower Preto River, upper Paraná River basin, Brazil. *Biota Neotropica*, vol. 17, no. 1. <http://dx.doi.org/10.1590/1676-0611-bn-2016-0196>.
- SANTOS, R.S., SILVA, J.P.C., COSTA, M.R. and ARAÚJO, F.G., 2015. O tamanho de primeira maturação como parâmetro para estabelecimento de tamanho mínimo de captura para corvina no Sudeste do Brasil. *Boletim do Instituto de Pesca*, vol. 41, no. 3, pp. 507-518.